


Map 路線圖



* Taxi & Minibus no. 26 are available to and from the front entrance of Hospital.
26

Bus

6, 15, 66	Central Bus Station
76	Causeway Bay
41A, 63	North Point
19	Siu Sai Wan

Minibus

5, 26	Causeway Bay
24M	Admiralty

Taxi

Available to and from the front entrance of Hospital

Parking

Parking is available for a fee for patients and visitors

Hong Kong Adventist Hospital – Stubbs Road

Address	40 Stubbs Road, Hong Kong 40
Telephone	(852) 3651 8888
E-mail	hkahinfo@hkah.org.hk
Website	www.hkah.org.hk

2018/12 Revised

Extending the Healing Ministry of Christ

Adventist 港
Health 安

Hong Kong Adventist Hospital • Stubbs Road

香港港安醫院•司徒拔道

Pediatric General Surgical Packages



It is natural for parents to be concerned about their child's health, particularly if the child requires surgery. Hong Kong Adventist Hospital is dedicated to providing comprehensive and professional care in a safe, friendly, and pleasant environment. In order to allow parents to assess the affordability of the necessary surgery, and to make a wise and stress free decision, we are pleased to provide the following surgical packages.

Circumcision

Circumcision is usually necessary because the foreskin is long, with a narrow opening preventing the patient from adequately cleaning the end of the penis. Benefits of circumcision include improved genital hygiene and comfort, as well as a reduced risk of infection.



Herniotomy

A hernia in childhood is due to a developmental abnormality or weakness that allows the bowel to push through the muscles, resulting in a visible bulge and occasionally strangulation of the bowel. Surgery involves returning the bowel to the abdominal cavity, and then repairing the muscles and tissues, so as to prevent any recurrences of the hernia.

Orchidopexy (unilateral)

This is a surgical procedure to place the testicle in its correct position in the scrotum. Failure of the testicle to descend into the scrotum by the time of birth is referred to as Cryptorchidism, and if not corrected may result in incomplete development and non-function of the testicle.



+ž

rhš

4	ž-L	T. Ý	6A,	}
ÑIĀg	A			
Ī				
ĪĪ				
6RĪ				
ÑIMĒ	I			
ĪĪ				
ĪĪ				
ĪĪĪ				

žGq
/RBe

žRĀ

leš

GžĪĪ

žRĀ

1h9i

PĪĪ

1h7hĀ

PĪĪ

1h'7hĪ

PĪĪ

+ CĪg

AMĒ

i

+ Īg

AMĒ

i R2

+ RBĪTĪ

+ žRB

Ī

PvRztnĪĪĪ

PvRztnRdbĪRdĪĪ

ĪĪ

+ Ī9 PĪĪK

d/ĪĪV

2ĪĪĪNĪKNDKRUJKNe

Ī

ĪĪĪe

PRE ĪĪ 1/2

2018/12 3i

