

Composition and Features of APS 自體血蛋白溶液的成份和特點

- Anti-inflammatory cytokines: Effectively counteracts joint inflammation
抗炎因子: 能有效對抗關節發炎
- Growth factors: Support cartilage protection and repair
生長因子: 有助於軟骨保護與修復
- Supraphysiological protein level: Higher than natural blood concentrations enhance therapeutic effects
抗炎及生長因子蛋白濃度比血液為高, 增強治療效果

Treatment Mechanism 治療機制

The prepared APS solution is injected directly into the affected joint, where it:

- 1 Restores protein balance
Suppresses excessive inflammation
調節關節內的蛋白平衡, 有效抑制炎症反應
- 2 Protects cartilage
Growth factors help slow degeneration and promote healing
發揮生長因子的保護作用, 促進軟骨修復
- 3 Reduces pain
By calming inflammation
通過消炎減輕膝關節疼痛
- 4 Enhances mobility
Patients experience improved knee function and flexibility
改善活動功能, 患者的膝關節功能和靈活性得到改善

As a completely autologous (self-derived) treatment, APS offers a natural, safe, and effective solution for managing knee osteoarthritis pain and preserving joint health. 由於這種療法完全採用患者自身的血液成分, 具有高度的生物相容性, 為退化性膝關節炎患者提供了一個安全有效的治療選擇。

Advantages 優勢



Day surgery procedure
Short hospital stay for few hours only
治療以日間手術形式進行
留院時間短至數小時



Performed under sterile conditions
Ensures safety and minimizes infection risk
整個過程在嚴格無菌條件下完成
確保安全性及減低感染風險



Minimal downtime
Most patients can resume light activities shortly after treatment
恢復期短
絕大部份患者在治療後不久即可恢復輕度活動

Patients who are appropriate for APS injection 適合自體血蛋白溶液注射治療的患者

Any patient suffered from knee osteoarthritis is appropriate for the APS injection, especially:

Mild to moderate degeneration patients
Studies have shown that APS is most effective in pain reduction in patient with early and moderate stage of osteoarthritis

輕微至中度退化性關節炎患者

研究顯示自體血蛋白溶液注射治療於早期至中期關節退化階段使用, 痛緩解效果更顯著

APS has received the following regulatory approvals/ registration: 自體血蛋白溶液 已獲取以下認證/註冊:

- CE Mark in Europe, the Middle East, and Africa (EMEA) on 2012
- FDA clearance as a Platelet-rich plasma (PRP) product on 2023
- In the Asia Pacific region, APS has been cleared by local health authority as treatment of degenerative knee osteoarthritis in Australia, New Zealand, Japan, and Singapore
- Introduced to Hong Kong and Macau in 2016 and 2018 respectively, registered in Hong Kong Department of Health in 2020 as Medical Device [HKMD No.200424]
- 2012年取得歐洲、中東及非洲地區的CE標誌
- 2023年獲美國食品藥物管理局(FDA)核准作為PRP (高濃度血小板血漿)療法
- 亞太地區: 獲澳洲、紐西蘭、日本及新加坡衛生機構批准作為療法治療退化性膝關節炎
- 香港及澳門: 分別於2016年及2018年開始引入, 2020年於香港衛生署註冊為醫療器械 [HKMD No.200424]


FAQ 常見問題

1. **Can APS cure my Knee Osteoarthritis?**
Currently, there is no medical evidence to support any single therapy that can completely cure degenerative knee osteoarthritis. However, APS injections can slow progression and reduce pain. Early treatment yields better results.
自體血蛋白溶液注射治療能治癒我的退化性膝關節炎嗎?
現時並沒有醫學研究指出任何一種治療能夠徹底治癒退化性膝關節炎。自體血蛋白溶液注射治療有效減緩退化和減輕疼痛, 越早治療效果越好。

2. **How long does it take for APS injection to be effective?**
Clinical studies showed that it will take about 4-6 weeks for patient to feel significant pain reduction.
APS注射治療需要多久才能見效?
研究顯示患者會於4至6星期感到疼痛顯著減少。

3. **Do I need repetitive injection?**
Most patients experience effective pain relief for 1 to 2 years after a single injection (Effect might vary and depends on the knee condition). When the therapeutic effects gradually diminish, patients can receive additional injections based on their needs.
是否需要重複注射?
大多數患者在接受單次注射後, 痛緩解1至2年(視乎患者情況而定), 當療效逐漸減弱時, 患者可依據實際需求再次接受注射治療。

MAP 路線圖



Hong Kong Adventist Hospital – Stubbs Road
香港港安醫院 — 司徒拔道

④ 40 Stubbs Road, Hong Kong
香港司徒拔道40號

📞 (852) 3651 8990 📩 hkahinfo@hkah.org.hk



Facebook



Make an Appointment



More Related Information and Video
更多相關資訊及影片

Contact us today to learn more or schedule an appointment.
請立即聯絡我們以了解更多資訊或預約服務。

Clinical Studies Reference: 臨床醫學文獻參考:

1. King W, van der Weegen W, Van Drunen R, Soons H, Toler K, Woodell-May J. White blood cell concentration correlates with increased concentrations of IL-1ra and improvement in WOMAC pain scores in an open-label safety study of autologous protein solution. *J Exp Orthop.* 2016;3:9. 2. Kon E, Engebretsen L, Verdonk P, Nehrer S, Filardo G. Clinical outcomes of knee osteoarthritis treated with an autologous protein solution injection: A 1-year pilot double-blinded randomized controlled trial. *J Exp Orthop.* 2016;3:9. 3. Kon E, Engebretsen L, Verdonk P, Nehrer S, Filardo G. Autologous protein solution injections for the treatment of knee osteoarthritis: 3-year results. *J Exp Orthop.* 2023;10(1):9. doi:10.1186/s40634-023-00501-5. 4. Kon E, Verdonk P, Engebretsen L, Nehrer S, Di Martino A, Filardo G. Duration effects of a single intra-articular injection of Autologous Protein Solution (APS) in patients with knee osteoarthritis (OA). 15th World Congress of the International Cartilage Repair Society, Vancouver, Canada, Program #12.4.4 (#437), Oct. 5-8, 2019.

Hong Kong Adventist Hospital - Stubbs Road reserves the right to update the fee schedule, brochure, terms and conditions. Any change of fee schedule will be announced and notices will be published in advance, in accordance with the statutory notice period. Any other changes except the fee schedule may be made at any time as it sees fit without prior notice, you are advised to check the latest information before using our services. 香港港安醫院—司徒拔道保留修訂收費表、單張內容、條款及細則之權利。任何的收費表調整將會根據法定的通知期提前發出通告和作出宣布。其他與收費表以外的相關修訂, 則可能不作另行通知。閣下於使用本院服務前查閱最新資訊。

SR-ORTH-APS-2025001

Adventist Health
Hong Kong Adventist Hospital - Stubbs Road
香港港安醫院・司徒拔道



AUTOLOGOUS PROTEIN SOLUTION INJECTION

自體血蛋白溶液注射治療

Extending the Healing Ministry of Christ

延續基督的醫治大能




What is Knee Osteoarthritis?

甚麼是退化性膝關節炎？

Osteoarthritis is a degenerative disease caused by long-term wear of cartilage in the joints, leading to inflammation, deformation, and bone spur growth. It primarily affects weight-bearing joints like the knees, hips, and fingers, with the knee joint being the most commonly affected area.

退化性關節炎是一種因關節軟骨長期磨損而導致的退化性疾病，會出現關節發炎、變形和骨刺增生。它主要影響經常承重的關節，如膝蓋、髖關節和手指關節，以膝關節最為常見。

Symptom 症狀



As joints degenerate, cartilage softens or develops pits, causing mild inflammation and knee pain on stairs or inclines, along with soreness, redness, swelling, and tightness. In later stages, cartilage may wear away completely, leading to irregular joint surfaces and bone spurs. Patients experience stiffness, limited movement, and deformities like "O-shaped feet," which reduce range of motion and cause weakness, especially when climbing stairs or squatting.

隨著關節退化，軟骨會軟化或出現凹洞，導致輕度發炎，患者在上下樓梯或斜坡上會感到膝部疼痛，並伴隨痠痛、紅腫和緊繃感。後期，軟骨可能完全磨損，造成關節表面不平和形成骨刺。患者會感到僵硬、活動受限，甚至出現「O型腳」的變形，這會降低關節的活動幅度，並在上下樓梯或蹲跪時感到無力。

What is Autologous Protein Solution Injection (APS)?

甚麼是自體血蛋白溶液注射治療(APS)？

APS is an innovative treatment that utilizes the patient's own blood-derived proteins to alleviate pain caused by knee osteoarthritis. Since APS is prepared from the patient's own blood, it carries no risk of allergic reactions or immune rejection, making it a safe and targeted therapy for degenerative knee joint pain.

自體血蛋白溶液注射治療是一種利用患者自身血液中的蛋白成分來治療膝骨關節炎疼痛的創新療法。這種治療方法完全採用患者自身的生物活性蛋白，因此不會引發過敏或排斥反應，特別適合用於退化性膝關節炎引起的疼痛治療。



How It Works 原理

The joints contain two key types of proteins:
人體關節內存在兩類關鍵蛋白：

Anti-inflammatory Cytokines ('Good' Protein) 抗炎因子（優質蛋白）

Help suppress inflammation and protect joint tissues
具有抑制炎症、保護關節的作用

Pro-inflammatory Cytokines ('Bad' Protein) 促炎因子（劣質蛋白）

Promote inflammation, leading to cartilage damage
會加劇炎症反應，導致軟骨損傷

製造軟骨



破壞軟骨

In a healthy joint, these proteins remain in balance. However, in osteoarthritis, pro-inflammatory cytokines increase, disrupting this equilibrium and accelerating cartilage breakdown.

在健康狀態下，這兩類蛋白維持著動態平衡。但當關節退化時，促炎因子會過度增加，破壞這種平衡。

Treatment Procedure 治療流程

STEP 1 程序 1

Blood Collection 抽血

- Draw approximately 60 mL of blood from a vein in your arm.
從手臂靜脈抽取約 60 毫升的血液
- The process is quick and safe, similar to a routine blood test.
整個抽血過程安全快速，類似常規血液檢查

STEP 2 程序 2

APS Preparation 製備

- The collected blood is immediately transferred to a sterile processing device.
抽取的血液會立即轉移至無菌分離器中
- Using a specialized centrifuge, the blood is processed and concentrated into a high-potency autologous protein solution (APS).
透過專用離心機進行分離與濃縮處理
- This step takes about 25 minutes.
約 25 分鐘即可完成



STEP 3 程序 3

Injection 注射

- The prepared APS solution (approximately 3-4 mL) is injected into the affected knee joint by a physician.
醫生將製備好約 3-4 毫升自體血蛋白溶液注射至受影響的膝關節內
- The injection is performed with precision to maximize therapeutic effects.
注射過程精準執行，以達到最佳治療效果



STEP 4 程序 4

Post-Treatment Observation 術後觀察

- After the injection, rest for about 1 hour for monitoring.
注射完成后需休息观察約 1 小時



Autologous Protein Solution (APS) Injection Therapy Package

4 Autologous Protein Solution (APS) Injection Therapy Package



Package Includes

Doctor procedure fee

Hospital inpatient care

Anesthesiologist fee

Operating room charges including equipment, consumables and instrument

Basic consumables or materials charges

Accommodation charge for Standard Ward

Package Charge(s)	Uilateral	Bilateral
HK\$	\$27,600	\$44,000

Important Information

1. The above packages are only valid for day cases in the Standard Ward.
2. The above packages do not meals during hospital stay.
3. Extra charges may apply under the following circumstances. Please contact our hospital staff for more details:
 - + Additional examinations or treatments not included in the package are required
 - + Any lab test, treatment, medication, procedures fee related to pre-existing condition
 - + Additional medications or medical supplies not included in the package are required
 - + Emergency services beyond the scheduled procedure are required.
 - + Complications arise during the procedure
 - + Changing bed within the same grade of accommodation
4. Eligibility for the above packages is based on the recommendation of a HKAH – SR specialist and each patient's actual condition. HKAH-SR reserves the right of final decision regarding the suitability of the packages for each patients.
5. Hong Kong Adventist Hospital - Stubbs Road reserves the right to update the fee schedule, brochure, terms and conditions. Any change of fee schedule will be announced and notices will be published in advance, in accordance with the statutory notice period. Any other changes except the fee schedule may be made at any time as it sees fit without prior notice, you are advised to check the latest information before using our services.

Appointments or Enquiries



Tel: (852) 3651 8629



WhatsApp: (852) 6463 0638

自體血蛋白溶液注射治療套餐

4 自體血蛋白溶液注射治療套餐

4 套餐包括

醫生手術費

醫生巡房費

麻醉科醫生收費

手術室費用，包括設備、消耗品和儀器

基本耗材或材料費用

普通房住宿收費

套餐收費	單側	雙側
港幣	\$27,600	\$44,000

注意事項

1. 以上費用只適用於普通房。
2. 以上套餐不包括住院期間的餐膳。
3. 以下情況將收取額外費用，詳情請向本院職員查詢：
 - + 計劃以外的檢查、治療
 - + 與既往病史相關的任何化驗、治療、藥物、費用
 - + 使用計劃以外的藥物和醫療物品
 - + 一般程序以外的緊急治療
 - + 過程中出現併發症
 - + 同等級病房內的換床費
4. 須視乎本院專科醫生的建議及求診者的實際情況而決定是否適合參與以上套餐。本院保留病人是否適用套餐的最終決定權。
5. 香港港安醫院一司徒拔道保留修訂收費表、單張內容、條款及細則之權利。任何的收費表調整將會根據法定的通知期提前發出通告和作出宣布。其他與收費表以外的相關修訂，則可能不作另行通知，閣下於使用本院服務前查閱最新資訊。

預約及查詢

 電話: (852) 3651 8629

 WhatsApp: (852) 6463 0638